





*Global Biogeochemical Cycles*

Supporting Information for

**Blue carbon storage capacity of eelgrass (*Zostera marina*) meadows: a global survey**

[Maria Emilia Röhr<sup>1,2\*</sup>, Marianne Holmer<sup>2</sup>, Julia K. Baum<sup>3</sup>, Mats Björk<sup>4</sup>, Diana Chin<sup>5</sup>, Lia Chalifour<sup>3</sup>, Stephanie Cimon<sup>6</sup>, Mathieu Cusson<sup>6</sup>, Martin Dahl<sup>4</sup>, Diana Deyanova<sup>4,7</sup>, J. Emmett Duffy<sup>8,\*\*</sup>, Johan S. Eklöf<sup>4</sup>, Julie K. Geyer<sup>9</sup>, John N. Griffin<sup>10</sup>, Martin Gullström<sup>6</sup>, Clara M. Hereu<sup>11</sup>, Masakazu Hori<sup>12</sup>, Kevin A Hovel<sup>13,\*\*</sup>, A. Randall Hughes<sup>14</sup>, Pablo Jorgensen<sup>15</sup>, Stephanie Kiriakopolos<sup>16,27</sup>, Per-Olav Moksnes<sup>17</sup>, Masahiro Nakaoaka<sup>18</sup>, Mary I. O'Connor<sup>19</sup>, Bradley Peterson<sup>4</sup>, Katrin Reiss<sup>20</sup>, Pamela L. Reynolds<sup>21,\*\*</sup>, Francesca Rossi<sup>22</sup>, Jennifer Ruesink<sup>23</sup>, Rui Santos<sup>24</sup>, John J. Stachowicz<sup>21</sup>, Fiona Tomas<sup>25,26</sup>, Kun-Seop Lee<sup>27</sup>, Richard K.F. Unsworth<sup>10</sup>, Christoffer Boström<sup>1</sup>

<sup>1</sup>Åbo Akademi University, Faculty of Science and Engineering, Environmental and Marine Biology, Artillerigatan 6, 20520 Åbo, Finland

<sup>2</sup>University of Southern Denmark, Department of Biology, Campusvej 55, 5230 Odense M, Denmark

<sup>3</sup>Department of Biology, University of Victoria, PO Box 1700 Station CSC, Victoria, British Columbia, V8W 2Y2, Canada.

<sup>4</sup>Department of Ecology, Environment and Plant Sciences, Stockholm University, 106 91 Stockholm, Sweden

<sup>5</sup>School of Marine and Atmospheric Sciences, Stony Brook University, 239 Montauk Highway, Southampton, NY 11968, USA

<sup>6</sup>Département des sciences fondamentales & Québec-Océan, Université du Québec à Chicoutimi, 555 boulevard de l'Université, Chicoutimi, Québec, G7H 2B1, Canada

<sup>7</sup>Institute for Biodiversity and Ecosystem Research at the Bulgarian Academy of Sciences, Sofia, Bulgaria

<sup>8</sup>Tennenbaum Marine Observatories Network, Smithsonian Institution, 647 Contees Wharf Road, Edgewater, MD 21037, USA

<sup>9</sup>Institute of Marine Sciences, University of North Carolina at Chapel Hill, Morehead City, NC, 28557, USA

<sup>10</sup>College of Science, Singleton Park, Swansea University, Swansea, SA28PP, UK

<sup>11</sup>Facultad de Ciencias Marinas, Universidad Autónoma de Baja California, Ensenada, Baja California, Mexico

<sup>12</sup>National Research Institute of Fisheries and Environment of Inland Sea, Fisheries Research and Education Agency, Hatsukaichi, Hiroshima 739-0452, Japan

<sup>13</sup>Department of Biology and Coastal & Marine Institute, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182

<sup>14</sup>Department of Marine and Environmental Science, Northeastern University, 430 Nahant Rd, Nahant, MA 01908 USA

<sup>15</sup>Geomare, Ensenada, Baja California, Mexico

<sup>16</sup>Estuary and Ocean Science Center and Department of Biology, San Francisco State University, Tiburon, CA, United States

<sup>17</sup>Department of Marine Sciences, University of Gothenburg, SE-405 30 Goteborg, Sweden

<sup>18</sup>Akkeshi Marine Station, Field Science Center for Northern Biosphere, Hokkaido University, 1 Aikappu, Akkeshi, Hokkaido 088-1113, Japan

<sup>19</sup>Department of Zoology and Biodiversity Research Centre, University of British Columbia, Vancouver, BC Canada V6T 1Z4

<sup>20</sup>Nord University, Faculty for Biosciences and Aquaculture, Postbox 1490, 8049 Bodø, Norway

<sup>21</sup>University of California, Davis, Data Science Initiative, One Shields Avenue, Davis, CA 95616.

<sup>22</sup>CNRS, Centre for marine Biodiversity, Exploitation and Conservation (MARBEF), University of Montpellier, PI E Bataillon, 34095 Montpellier

<sup>23</sup>Department of Biology, University of Washington, Seattle, WA, USA

<sup>24</sup>ALGAE -Marine Ecology Research Group, CCMar - Center of Marine Sciences, University of Algarve, Faro, Portugal

<sup>25</sup>IMEDEA, UIB-CSIC, Esporles, Islas Baleares, Spain

<sup>26</sup>Department of Fisheries and Wildlife, Oregon State University, Corvallis, OR, USA<sup>27</sup>Department of Biological Sciences, Pusan National University, Pusan, Korea

## **Contents of this file**

- Figures S1 to S2
- Tables S1 to S3




Figure S1. The relative contribution of phytoplankton to the  $\delta^{13}\text{C}$  of the sediment surface layer (0-5 cm) at the sites. Box plots represents first and third quartiles and are shown with medians (horizontal line), means (+). The whiskers represent the 2.5-97.5 percentiles.




Figure S2. Partial least square (PLS) regression model coefficient plot showing the relative importance of different predictor variables in the different ocean margin/seas. (Mud= mud content (%), SedDen= sediment density, salinity, sorting= degree of sorting ( $\phi$ ),  $^{15}\text{N}$  leaves=  $\delta^{15}\text{N}$  in *Z. marina* leaves, Zm Contr= *Z. marina* contribution to the sediment surface  $C_{org}$  pool (%), ShoDens= shoot density (shoots  $\text{m}^{-2}$ ), Ag biomass= aboveground biomass ( $\text{gDW m}^{-2}$ ), PON leaves= particulate organic nitrogen content of *Z. marina* leaves (%) and lat= latitude, ( $\text{gDW m}^{-2}$ ) on *Z. marina*  $C_{org}$  stocks. Predictor variables are ranked in order of importance (from the left to the right). Brown bars represent sediment characteristics, green bars represent seagrass-associated variables and blue bars environmental variables.

Table S1. Summary of site-specific sediment and environmental variables measured in this study. Country, location, abbreviation (Abb.), latitude (Lat.), longitude (Long.) in WGS 84 format, water depth (m), mean water temperature (°C), salinity (Sal.), mud content in the sediment surface (%), sediment dry density in the sediment surface (Dry dens. g cm<sup>-3</sup>), sediment organic carbon content in the sediment surface (SedOC, % DW), δ<sup>13</sup>C sediment surface, *Z. marina* contribution to the sediment surface C<sub>org</sub> pool (*Z. marina* contrib.(%)), degree of sorting (DS (φ)) and ocean margin/sea (WA= Western Atlantic, EA= Eastern Atlantic, WP= Western Pacific, EP= Eastern Pacific, MED= Mediterranean Sea, KS= Kattegat-Skagerrak , BS= Baltic Sea, BLS= Black Sea) at the sampling sites. SEM (n=3) is given.

Country	Location	Abb.	Lat.	Long.	Water depth (m)	Mean water temp. (°C)	Sal.	Mud content (%)	Dry dens. (g cm <sup>-3</sup> )	SedOC (% DW)
CAN	Tsawwassen	BC	49.00	-123.10	0.50	8.6	26.0	14.0±0.7	1.35±0.04	0.28±0.10
CAN	Prince Rupert	PR	54.19	-130.24	1.04	10.6	25.1	11.5±4.2	1.42±0.09	0.41±0.06
CAN	Baie-St-Ludger	QU	49.11	-68.17	1.75	8.0	21.0	26.6±2.2	1.42±0.09	0.45±0.08
USA	Bodega Bay	BB	38.32	-123.05	1.00	13.4	33.0	30.3±0.3	1.62±0.03	0.08±0.04
USA	Shinnecock Bay	LI	40.86	-72.45	0.54	12.3	30.3	15.5±5.2	1.30±0.02	0.27±0.07
USA	Dorothy Cove	MA	42.42	-70.91	2.00	14.0	32.4	34.2±4.3	1.46±0.04	0.29±0.04
USA	Yaquina Bay	OR	44.61	-124.01	0.15	12.3	36.0	15.6±2.9	2.04±0.02	0.62±0.04
USA	Coronado	SD	32.70	-117.17	1.82	18.5	35.0	7.6±2.1	1.28±0.11	0.22±0.03
USA	Point San Pablo	SF	37.95	-122.41	0.50	13.0	27.0	19.9±4.3	1.12±0.05	0.31±0.08
USA	Goodwin Islands	VA	37.22	-76.40	1.00	19.7	18.3	9.0±3.6	1.43±0.02	0.17±0.05
USA	Willapa Bay	WA	46.47	-124.02	3.16	12.6	23.0	36.9±1.8	0.93±0.07	0.77±0.10
USA	Shackleford Island	NC	34.67	-76.57	0.65	12.6	25.0	21.5±3.7	1.68±0.02	0.36±0.02
MEX	San Quintin Bay	MX	30.42	-115.96	1.15	17.6	34.0	24.3±16.7	1.45±0.06	0.55±0.07
JPN	Akkeshi-ko	JN	43.02	144.90	1.50	8.5	26.0	60.3±4.8	1.00±0.07	0.89±0.01
JPN	Ikunoshima	JS	34.30	132.91	2.00	15.3	32.5	75.9±5.6	0.95±0.09	1.22±0.11
KOR	Dongdae Bay	KO	34.89	128.02	0.40	16.5	31.3	81.3±0.5	1.02±0.05	1.28±0.06
FRA	Bouzigues	FR	43.45	03.66	2.10	16.5	38.8	76.3±0.7	1.33±0.21	2.34±0.06
UK	Porth Dinllaen	UK	52.94	-04.56	2.00	11.5	34.4	43.6±5.1	1.33±0.13	0.44±0.12
POR	Culatra	CU	37.00	07.83	1.18	18.1	35.3	25.0±1.6	1.20±0.05	1.65±0.15
BUL	Garden	GA	42.43	27.65	5.00	15.7	16.5	2.3±0.4	1.28±0.01	2.35±0.14
BUL	Ropotamo	RO	42.33	27.75	2.50	15.7	16.5	7.5±2.9	1.26±0.03	4.68±0.37
NOR	Rövika	NO	67.21	15.00	1.00	7.3	18.5	3.1±0.2	1.51±0.01	0.12±0.01
SWE	Snäckebackebukten	SN	58.36	11.56	0.20	10.0	24.0	40.4±8.2	0.51±0.08	5.37±0.86
SWE	Torgestad	TO	58.34	11.55	1.70	10.0	24.0	32.2±0.8	1.46±0.09	0.41±0.14
SWE	Lindholmen	LN	58.26	11.48	1.60	10.0	24.0	55.6±6.9	0.22±0.02	10.65±0.12
SWE	Bökevik	BO	58.25	11.45	1.60	10.0	24.0	61.1±7.6	0.89±0.09	0.66±0.09
SWE	Hakefjord	HA	58.04	11.80	1.60	10.0	24.0	61.0±17.3	1.37±0.06	0.86±0.11
SWE	Wallhamn	WL	58.01	11.71	1.10	10.0	24.0	29.7±23.1	0.59±0.06	3.96±0.54
SWE	Storebrorn	ST	57.89	11.66	2.50	10.0	24.0	27.6±5.6	0.45±0.05	2.44±0.15
SWE	Finnsbo	FI	58.30	11.78	1.70	10.0	24.0	68.1±13.5	0.42±0.05	3.78±0.75
SWE	Kristineberg	KR	58.25	11.45	2.00	10.0	24.0	52.0±13.5	0.96±0.15	2.61±0.05
SWE	Stora Sand	SS	58.81	11.66	3.80	8.8	6.4	2.6±0.4	1.21±0.02	0.26±0.05
SWE	Långskär	LS	58.80	11.70	2.30	8.8	6.4	4.3±0.7	1.50±0.07	0.45±0.08
SWE	Torö	TR	58.81	11.65	3.50	8.8	6.4	5.4±1.0	1.42±0.11	0.42±0.02
DEN	Nyborg	NY	55.30	10.83	2.50	10.0	20.0	0.5±0.3	1.17±0.03	0.10±0.01
DEN	Kertinge	KE	55.44	10.55	2.50	10.0	20.0	27.1±1.5	1.15±0.05	3.23±0.24
DEN	Løgstør	LØ	57.01	09.05	2.50	10.0	20.0	4.0±0.4	1.23±0.03	0.31±0.09
DEN	Lovns	LO	56.62	09.28	2.50	10.0	20.0	17.8±3.0	1.22±0.09	1.53±0.09
DEN	Agero 3	AG3	56.67	08.58	2.50	10.0	20.0	30.4±6.5	1.24±0.09	2.30±0.08
DEN	Visby	VI	56.77	08.48	2.50	10.0	20.0	23.0±4.2	1.25±0.02	2.18±0.20
DEN	Lunkebugt	LU	54.99	10.65	2.50	10.0	20.0	33.2±7.4	1.23±0.23	1.71±0.81
DEN	Thurøbund	TH	55.04	10.69	2.50	10.0	20.0	34.7±2.8	1.27±0.03	5.78±0.51
DEN	Agero 12	AG12	56.69	08.58	2.50	10.0	20.0	27.9±8.0	1.35±0.17	0.29±0.14
DEN	Dalby	DA	55.53	10.60	2.50	10.0	20.0	8.1±1.6	1.37±0.03	0.12±0.01
FIN	Sackholm	SA	60.12	21.86	2.50	10.0	6.5	12.5±2.0	1.36±0.04	0.26±0.03
FIN	Ängsö	ÄN	60.11	21.71	2.50	10.0	6.5	4.2±2.1	1.36±0.05	0.20±0.01
FIN	Kolaviken	KL	59.82	22.99	2.50	10.0	6.5	1.9±0.2	1.34±0.04	0.13±0.01
FIN	Ryssholmen	RY	59.83	23.08	2.50	10.0	6.5	2.7±0.6	1.34±0.05	0.16±0.01
FIN	Tvärminne	TV	59.84	23.24	2.50	10.0	6.5	9.2±1.9	1.33±0.03	0.20±0.02
FIN	Fårö	FÅ	59.92	21.81	2.50	10.0	6.5	5.1±1.4	1.32±0.03	0.13±0.01
FIN	Lyddaren	LY	60.13	21.44	2.50	10.0	6.5	3.2±2.4	1.34±0.17	0.45±0.09
FIN	Långören	LÅ	59.88	21.74	2.50	10.0	6.5	2.9±2.1	1.42±0.05	0.19±0.02

FIN	Hummelskär	HU	60.03	21.71	2.50	10.0	6.5	$9.0 \pm 2.6$	$1.33 \pm 0.01$	$0.35 \pm 0.02$
FIN	Jänisholm	JÄ	60.21	21.70	2.50	10.0	6.5	$7.2 \pm 2.2$	$1.37 \pm 0.08$	$0.33 \pm 0.14$

Table S2. Summary of site-specific plant variables measured in this study. Country, location, abbreviation (Abb.), shoot density (shoots m<sup>-2</sup>), δ <sup>13</sup>C of *Z. marina* leaves, δ<sup>13</sup>C of *Z. marina* rhizomes, C:N ratio of *Z. marina* (C:N), *Z. marina* above -and belowground biomass (AB and BB, gDW m<sup>-2</sup>) and root:shoot- ratio (R:S) at the sampling sites. SEM (n=3) is given.

Country	Location	Abb.	Lat.	Long.	Water depth (m)	Mean water temp. (°C)	Sal.	Mud content (%)	Dry dens. (g cm <sup>-3</sup> )	SedOC (% DW)
CAN	Tsawwassen	BC	49.00	-123.10	0.50	8.6	26.0	14.0±0.7	1.35±0.04	0.28±0.10
CAN	Prince Rupert	PR	54.19	-130.24	1.04	10.6	25.1	11.5±4.2	1.42±0.09	0.41±0.06
CAN	Baie-St-Ludger	QU	49.11	-68.17	1.75	8.0	21.0	26.6±2.2	1.42±0.09	0.45±0.08
USA	Bodega Bay	BB	38.32	-123.05	1.00	13.4	33.0	30.3±0.3	1.62±0.03	0.08±0.04
USA	Shinnecock Bay	LI	40.86	-72.45	0.54	12.3	30.3	15.5±5.2	1.30±0.02	0.27±0.07
USA	Dorothy Cove	MA	42.42	-70.91	2.00	14.0	32.4	34.2±4.3	1.46±0.04	0.29±0.04
USA	Yaquina Bay	OR	44.61	-124.01	0.15	12.3	36.0	15.6±2.9	2.04±0.02	0.62±0.04
USA	Coronado	SD	32.70	-117.17	1.82	18.5	35.0	7.6±2.1	1.28±0.11	0.22±0.03
USA	Point San Pablo	SF	37.95	-122.41	0.50	13.0	27.0	19.9±4.3	1.12±0.05	0.31±0.08
USA	Goodwin Islands	VA	37.22	-76.40	1.00	19.7	18.3	9.0±3.6	1.43±0.02	0.17±0.05
USA	Willapa Bay	WA	46.47	-124.02	3.16	12.6	23.0	36.9±1.8	0.93±0.07	0.77±0.10
USA	Shackleford Island	NC	34.67	-76.57	0.65	12.6	25.0	21.5±3.7	1.68±0.02	0.36±0.02
MEX	San Quintin Bay	MX	30.42	-115.96	1.15	17.6	34.0	24.3±16.7	1.45±0.06	0.55±0.07
JPN	Akkeshi-ko	JN	43.02	144.90	1.50	8.5	26.0	60.3±4.8	1.00±0.07	0.89±0.01
JPN	Ikunoshima	JS	34.30	132.91	2.00	15.3	32.5	75.9±5.6	0.95±0.09	1.22±0.11
KOR	Dongdae Bay	KO	34.89	128.02	0.40	16.5	31.3	81.3±0.5	1.02±0.05	1.28±0.06
FRA	Bouzigues	FR	43.45	03.66	2.10	16.5	38.8	76.3±0.7	1.33±0.21	2.34±0.06
UK	Porth Dinllaen	UK	52.94	-04.56	2.00	11.5	34.4	43.6±5.1	1.33±0.13	0.44±0.12
POR	Culatra	CU	37.00	07.83	1.18	18.1	35.3	25.0±1.6	1.20±0.05	1.65±0.15
BUL	Garden	GA	42.43	27.65	5.00	15.7	16.5	2.3±0.4	1.28±0.01	2.35±0.14
BUL	Ropotamo	RO	42.33	27.75	2.50	15.7	16.5	7.5±2.9	1.26±0.03	4.68±0.37
NOR	Rövika	NO	67.21	15.00	1.00	7.3	18.5	3.1±0.2	1.51±0.01	0.12±0.01
SWE	Snäckebackebukten	SN	58.36	11.56	0.20	10.0	24.0	40.4±8.2	0.51±0.08	5.37±0.86
SWE	Torgestad	TO	58.34	11.55	1.70	10.0	24.0	32.2±0.8	1.46±0.09	0.41±0.14
SWE	Lindholmen	LN	58.26	11.48	1.60	10.0	24.0	55.6±6.9	0.22±0.02	10.65±0.12
SWE	Bökevik	BO	58.25	11.45	1.60	10.0	24.0	61.1±7.6	0.89±0.09	0.66±0.09
SWE	Hakefjord	HA	58.04	11.80	1.60	10.0	24.0	61.0±17.3	1.37±0.06	0.86±0.11
SWE	Wallhamn	WL	58.01	11.71	1.10	10.0	24.0	29.7±23.1	0.59±0.06	3.96±0.54
SWE	Storebrorn	ST	57.89	11.66	2.50	10.0	24.0	27.6±5.6	0.45±0.05	2.44±0.15
SWE	Finnsbo	FI	58.30	11.78	1.70	10.0	24.0	68.1±13.5	0.42±0.05	3.78±0.75
SWE	Kristineberg	KR	58.25	11.45	2.00	10.0	24.0	52.0±13.5	0.96±0.15	2.61±1.05
SWE	Stora Sand	SS	58.81	11.66	3.80	8.8	6.4	2.6±0.4	1.21±0.02	0.26±0.05
SWE	Långskär	LS	58.80	11.70	2.30	8.8	6.4	4.3±0.7	1.50±0.07	0.45±0.08
SWE	Torö	TR	58.81	11.65	3.50	8.8	6.4	5.4±1.0	1.42±0.11	0.42±0.02
DEN	Nyborg	NY	55.30	10.83	2.50	10.0	20.0	0.5±0.3	1.17±0.03	0.10±0.01
DEN	Kertinge	KE	55.44	10.55	2.50	10.0	20.0	27.1±1.5	1.15±0.05	3.23±0.24
DEN	Løgstør	LØ	57.01	09.05	2.50	10.0	20.0	4.0±0.4	1.23±0.03	0.31±0.09
DEN	Lovns	LO	56.62	09.28	2.50	10.0	20.0	17.8±3.0	1.22±0.09	1.53±0.09
DEN	Agero 3	AG3	56.67	08.58	2.50	10.0	20.0	30.4±6.5	1.24±0.09	2.30±0.08
DEN	Visby	VI	56.77	08.48	2.50	10.0	20.0	23.0±4.2	1.25±0.02	2.18±0.20
DEN	Lunkebugt	LU	54.99	10.65	2.50	10.0	20.0	33.2±7.4	1.23±0.23	1.71±0.81
DEN	Thurøbund	TH	55.04	10.69	2.50	10.0	20.0	34.7±2.8	1.27±0.03	5.78±0.51
DEN	Agero 12	AG12	56.69	08.58	2.50	10.0	20.0	27.9±8.0	1.35±0.17	0.29±0.14
DEN	Dalby	DA	55.53	10.60	2.50	10.0	20.0	8.1±1.6	1.37±0.03	0.12±0.01
FIN	Sackholm	SA	60.12	21.86	2.50	10.0	6.5	12.5±2.0	1.36±0.04	0.26±0.03
FIN	Ängsö	AN	60.11	21.71	2.50	10.0	6.5	4.2±2.1	1.36±0.05	0.20±0.01
FIN	Kolaviken	KL	59.82	22.99	2.50	10.0	6.5	1.9±0.2	1.34±0.04	0.13±0.01
FIN	Ryssholmen	RY	59.83	23.08	2.50	10.0	6.5	2.7±0.6	1.34±0.05	0.16±0.01
FIN	Tväminne	TV	59.84	23.24	2.50	10.0	6.5	9.2±1.9	1.33±0.03	0.20±0.02
FIN	Fårö	FÅ	59.92	21.81	2.50	10.0	6.5	5.1±1.4	1.32±0.03	0.13±0.01
FIN	Lyddaren	LY	60.13	21.44	2.50	10.0	6.5	3.2±2.4	1.34±0.17	0.45±0.09
FIN	Långören	LÅ	59.88	21.74	2.50	10.0	6.5	2.9±2.1	1.42±0.05	0.19±0.02
FIN	Hummelskär	HU	60.03	21.71	2.50	10.0	6.5	9.0±2.6	1.33±0.01	0.35±0.02
FIN	Jänisholm	JÄ	60.21	21.70	2.50	10.0	6.5	7.2±2.2	1.37±0.08	0.33±0.14

Table S3. The literature source used for plankton  $\delta^{13}\text{C}$  signal in the stable isotope mixing model analysis.

Site	$\delta^{13}\text{C}$	Country	Ocean margin/ Sea	Reference:
FR	-22.8	FR	Mediterranean Sea	Pernet et al. 2012
KO	-22.0	KO	Western Pacific	Keun-Kang et al. 2015
PR	-20.6	CA	Eastern Pacific	Goering et al. 1990
MA	-21.0	US	Western Atlantic	Tagliabue and Bopp 2008
NO	nd.	NO	Eastern Atlantic	nd.
OR	-22.0	US	Eastern Pacific	Conway-Cranos et al. 2015
LI	-21.0	US	Western Atlantic	Tagliabue and Bopp 2008
QU	-21.8	CA	Western Atlantic	Winkler and Cabrol <i>unpublished data</i>
SN	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
TO	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
LN	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
BO	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
HA	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
WL	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
ST	-22.5	SWE	Kattegat-Skagerrak	Tiselius and Fransson. 2015
VA	-21.0	US	Western Atlantic	Tagliabue and Bopp 2008
WA	-22.0	US	Eastern Atlantic	Conway-Cranos et al. 2015
JS	-21.9	JP	Western Pacific	Miyayima et al. 2015
SD	-20.7	US	Eastern Pacific	Jorgensen et al. 2007
MX	-20.7	MX	Eastern Pacific	Jorgensen et al. 2007
SF	-23.0	US	Eastern Pacific	Tagliabue and Bopp 2008
NC	-21.0	US	Western Atlantic	Tagliabue and Bopp 2008
JN	-18.9	JP	Western Pacific	Kajihara et al. 2010
BC	-20.6	CA	Eastern Pacific	Conway-Cranos et al. 2015
UK	-26.0	UK	Eastern Atlantic	Tagliabue and Bopp 2008
BB	-23.0	US	Eastern Pacific	Tagliabue and Bopp 2008
NY	-17.5	DK	Kattegat-Skagerrak	Röhr et al. 2016
KE	-17.5	DK	Kattegat-Skagerrak	Röhr et al. 2016
LØ	-16.4	DK	Kattegat-Skagerrak	Röhr et al. 2016
LO	-17.3	DK	Kattegat-Skagerrak	Röhr et al. 2016
AG3	-18.6	DK	Kattegat-Skagerrak	Röhr et al. 2016
AG12	-17.5	DK	Kattegat-Skagerrak	Röhr et al. 2016

<b>VI</b>	-17.4	DK	Kattegat-Skagerrak	Röhr et al. 2016
<b>LU</b>	-17.5	DK	Kattegat-Skagerrak	Röhr et al. 2016
<b>TH</b>	-17.5	DK	Kattegat-Skagerrak	Röhr et al. 2016
<b>DA</b>	-17.5	DK	Kattegat-Skagerrak	Röhr et al. 2016
<b>SA</b>	-16.4	FIN	Baltic Sea	Röhr et al. 2016
<b>ÄN</b>	-15.3	FIN	Baltic Sea	Röhr et al. 2016
<b>KL</b>	-20.2	FIN	Baltic Sea	Röhr et al. 2016
<b>RY</b>	-22.1	FIN	Baltic Sea	Röhr et al. 2016
<b>TV</b>	-23.0	FIN	Baltic Sea	Röhr et al. 2016
<b>FÅ</b>	-22.1	FIN	Baltic Sea	Röhr et al. 2016
<b>LY</b>	-22.3	FIN	Baltic Sea	Röhr et al. 2016
<b>LÅ</b>	-22.9	FIN	Baltic Sea	Röhr et al. 2016
<b>HU</b>	-24.3	FIN	Baltic Sea	Röhr et al. 2016
<b>JÄ</b>	-24.6	FIN	Baltic Sea	Röhr et al. 2016
<b>CU</b>	-16.6	PO	Eastern Atlantic	Santos et al. <i>unpublished data</i>
<b>FI</b>	-22.5	SWE	Kattegat- Skagerrak	Tiselius and Fransson. 2015
<b>GA</b>	-21.8	BU	Black Sea	Fry et al. 1991
<b>KR</b>	-22.5	SWE	Kattegat- Skagerrak	Tiselius and Fransson. 2015
<b>RO</b>	-21.8	BU	Black Sea	Fry et al. 1991
<b>SS</b>	-22.9	SWE	Baltic Sea	Röhr et al. 2016
<b>LS</b>	-22.9	SWE	Baltic Sea	Röhr et al. 2016
<b>TR</b>	-22.9	SWE	Baltic Sea	Röhr et al. 2016